



CGM/NMET/Lodhva block/650 や/24-25

Date: 1311 JUL 2024

To,
Director,
National Mineral Exploration Trust
Ministry of Mines, Government of India.
New Delhi, Delhi-110001.

Subject: Allocation of funds Rs 88.93 lakhs (Rupees eighty eight Lakhs ninty three Thousand) for G2 level Geological exploration for Limestone in Lodhva Block, Gir Somnath District, Gujarat State under NMET.

Respected Sir,

In the past, NMET has sanctioned several Geochemical analysis and Geological exploration projects proposed by CGM - Gujarat. These projects have been successfully executed by CGM - Gujarat. Currently, NMET has also provided funding for three ongoing geological exploration projects in the Kutch district.

The coastal region of Saurashtra, spanning from Okha in the west to Bhavnagar in the east, is a notable area in India characterized by the prevalent presence of Milliolite limestone. These formations extend up to 200 kilometers inland from the coastline. The entire coastal region of Saurashtra holds substantial potential for chemical grade and cement grade limestone deposits. Consequently, various private cement factories and Gujarat Alkalis have established soda ash plants in this area.

Currently, both cement and soda ash plants are encountering a shortage of raw limestone supply from the Saurashtra region. Consequently, they have resorted to importing raw limestone from other regions within the country and even overseas. Given the existing demand for limestone, it is crucial that we explore the remaining promising limestone reserves in the Saurashtra region to ensure efficient sourcing of raw materials from nearby locations.

Therefore, we are presenting a comprehensive proposal for conducting G2 level geological exploration for limestone in Lodhva Block with a total cost of Rs 88.93 lakhs funded by NMET. We would greatly appreciate your prompt approval of this proposal.

Thanking You.

For CGM - Gujarat

Dr. Dhaval Patel, IAS

Commissioner of Geology and mining. Gujarat State, Gandhinagar

Enclosed: As above

Copy to: (1) DDG - SU, GSI - Gujarat.

(2) Joint Secretory, Industries and Mines department, New Sachivalaya, Gandhinagar.

Proposal for Lodhva Block, Gir Somanth District, Gujarat State for G2 Stage Mineral Exploration under NMET



Commodity: Limestone

Ву

Commissioner of Geology and Mining Gujarat

Place: Gandhinagar Date: 29 JULY 2024

Summary of the Block for G2 stage exploration

	Features	Details						
	Block ID	CGM/NMET/I	limestone/06/20	24-25				
	Current Exploration Agency	Commissione	Miliolite Formation					
	Previous Exploration Agency	Commissione						
	Commodity	Limestone						
	Mineral Belt	Miliolite Form						
	Completion Period with entire Time schedule to complete the project	06 months						
	Objectives	To assess & identify qualitative quantitative mineral resource of Limes mineral at G2 stage in the proposed larea.						
	Whether the work will be carried out by the proposed agency or through outsourcing and details thereof.							
	Name/ Number of Geoscientists	Geologist: 1 (Field) + 1(HQ)						
	Expected Field days (Geology, Geophysics, Surveyor)	30 days: Field Geologist 40 days: HQ Geologist 20 days: Surveyor						
1.	Location		92.					
	Co-ordinates (Latitude, Longitude) of Block Boundary	Block corner points	Latitude	Longitude				
		1	20°51'13.08"N	70°35'6.65"E				
		2	20°51'16.47"N	70°35'58.28"E				
		3	20°50'33.01"N	70°36'40.89"E				
		4	20°50'29.12"N	70°35'1.70"E				
	Villages	Lodhva						
	Tehsil/ Taluk	Sutrapada						
	District	Gir Somnath						
	State	Gujarat		-				
2.	Area (hectares/ square kilometres)							
	Block Area	300 hectares						
	Forest Area	Nil						
	Government Land Area	NA						
	Private Land Area	NA .						
3.	Accessibility							
	Nearest Rail Head	Kodinar railway station – 9.00 km						
	Road	NH 101 - 1.5	km					
	Airport	Diu airport -	34.50 km					

Block Id: CGM/NMET/Limestone/06/2024-25 Commissioner of Geology and Mining, Gujarat

4.	Hydrography	
i i	Local Surface Drainage Pattern (Channels)	Sub-parallel Pattern
1	Rivers/ Streams	Small nallas presents in the area.
5.	Climate	
3	Mean Annual Rainfall	824 mm
9	Temperatures (December) (Minimum) Temperatures (June) (Maximum)	Minimum – 16° C Maximum – 36° C
6.	Topography	
3	Toposheet Number	41L/9
9	Morphology of the Area	The proposed area is generally covered by dr barren and agricultural fields. The topography of the proposed region is looking gently slop toward west direction. The elevation of the are ranges from 30mts to 52mts above mean selevel.
2-10-1 (5)	Availability of baseline geoscience data	
	Geological Map	Plate-1
- 1	Geochemical Map	Not available
1	Geophysical Map (Aerogeophysical, Ground geophysical, Regional as well as local scale GP maps)	Not available
	Justification for taking up G2 stage mineral exploration	 The region was primarily investigated by CGM in the year 1965 and 1995 to 1997 by means of Geological mapping using Remote sensing techniques, surface exposure mapping and sample collection of the proposed area. The recommendation of this report indicates Coastal areas of Saurashtra region have milliolitic limestone deposit these deposits have very less overburded and occurred in form of low ridges and low mounds. So, it can be easily minable deposit. Also, other facilities like Roads and transport, Labour, etc. are easily available in this region. Additionally, In the year 1979 to 1981 CGN had carried out core drilling in the 900mt away in eastern direction from the proposed block and borehole lithologindicated High-grade limestone in the region.

We have collected 08 grab samples during the recent geological travers survey of CGM / GMRDS geologist, the analysis results of these samples indicated CaO% from 47.87% to 53.43% (Refer Annexure - 1)
 The proposed block is surrounded by the existing working Limestone leases, so this area can be studied as an extension of the already existing mineable deposits. Thus, the block is suggested for G2 level of exploration.

Detailed description:

1. Block Summary

Physiography

The study area is located in the Sutrapada taluka, Gir Somnath district in the western coastal region of Gujarat. The area under investigation is a coastal plain, where major portion is flat agricultural fields leaving patches of limestone out crops in form of small low mounds and ridges more or less parallel to coast line. The limestone ridges are practically covering the sea-coast line. The lying slope of low-lying ridges and mounds generally merges into coastal plain. The rivers Hiran and Saraswati, both perennial flow through the area and are fed by numerous streams all originating in the highlands to the north.

Background Geology (Regional Geology & Geology of the Block).

The three major lithounits, i.e. Gaj, Dwarka and Miliolite Formations occur broadly in three linear belts parallel to the coastline. Chaya Formation, restricted near the coastline, is porous, semi-consolidated limestone rich in fossil fragments. The exposures of Gaj Formation are present farthest from the coast near the exposed Deccan Trap hillocks and Miliolite Formation along the coastline whereas the Dwarka Formation is exposed in between. The Miliolite Formation has covered all the pre-existing lithounits.

The southern coast of Saurashtra from Okha in the west to Bhavnagar in the east is unique in India, being the only place where nearly pure carbonate sedimentation has taken place during the quaternary period under warm shallow and agitating water. These limestones are known as Milliolite limestone. These occurrences are also observed inland upto 200 kms from the coastal line.

The general stratigraphic succession of the region is as follows:

Stratigraphic Unit	Lithology	Age
Recent Deposits	Coastal Sea Sand dunes, soil and alluvium	Holocene
Chhaya Formation	Shelly limestone, Coraline limestone	Holocene to late Pleistocene
Milliolite Formation	Limestones with shell of Milolina	Pleistocene
Dwarka Formation	Fossiliferous Limestone	Pliocene
Gaj Formation	Alternate sequence of clays and limestone	Miocene
Deccan Trap	Basalt dolerite dykes	Cretaceous to Eocene

Block Id: CGM/NMET/Limestone/06/2024-25 Commissioner of Geology and Mining, Gujarat

Gai Formation

Gaj formation includes an intercalated sequence of marl/clay and limestone, dominated by the former. The marl/clay is light yellow to grey and it is difficult to distinguish between the two in transitional cases. Gaj limestone is light to deep yellow to brown, compact, and cryptocrystalline. At places, this limestone contains very little free silica or is almost devoid of it. The thickness of limestone band is highly variable and often a quite thick band pinches abruptly within short distance., In almost all places, where Gaj Formation is exposed a, thin crust (less than 2 m thick) of compact, dark coloured limestone, which is a secondary leaching product, is present followed downward by Gaj clay/marl. Gaj Formation is rich in marine mega- and micro- fossils. Mega fossils include pelecypods, gastropods, cephalopods, echinoids, alcyonaria, decapoda and bryozoa. Earlier workers have assigned an Upper Miocene age to Gaj Formation but Mathur et al (1980) advocated that the Faj Formation is at least of Mid Miocene age, if not older, as it contains fossils of Taberina malabarica a Lower to Middle Miocene foraminifer.

Dwarka Formation

Dwarka Formation is equivalent to the Dwarka bed of Fedden (1884). It overlies the Gaj Formation. Although there are pebbly or conglomeratic bands at the contacts, in most places the relation is conformable. The Dwarka Formation, as exposed in this area, generally comprises grey and gritty to sandy limestone almost devoid of megafossils. Near Dwarka, the Dwarka formation was divided into Upper and Lower Dwarka Member by Jain and Agarwal (1989-90). The lower Member, which overlies Gaj Formation, is an intercalated sequence of clay and kankar, friable/compact sandstone, yellow and red marl beds and yellow ferruginous and grayish sandy limestone. The Upper Dwarka is mainly bioclastic limestone, often cavernous and contains very little silica. In the mapped area the exposed Dwarka Formation is mostly represented by Lower Dwarka Member of Jain and Agarwal (1990), although there are some pockets, which are less in free silica.

Milliolite Limestone

The consolidated Quaternary deposits are termed as Miliolite Formation because of preponderance of tests of foraminifer Miliolidae. This is the most widespread lithounit in the coastal belt overlapping in all earlier lithounits. The colour of the limestone varies from white to pale yellow to pink. In majority of places the miliolitic limestone varies from white to pale yellow, relatively soft, friable, thinly laminated with alternate bands rich in foraminiferal tests and micrite. The Milliolite occur as whitish, buff coloured, current bedded limestone in the form of coastal ridge This Limestone primarily comprise of broken shells of foraminifer-milioline around which calcite grains have been formed. However, this is not a pure limestone and thin horizons of sandstone, grit and conglomerate are also found within Milliolites. The

Milliolite are thought to be windblown deposits and are also found occurring along the hill slopes further inland.

Chhaya Formation

This is a semi-consolidated highly porous limestone occurring near the present day coast line and rich in both mega and micro-fossils. The thickness shows wide variation decreasing away from the coast. It is generally a bit ferruginous particularly below the ground level.

Soil and Alluvium

Major portion of the area is covered with soil or alluvium. This can be divided as the type formed due to weathering of limestone and the type formed due to deposition of clays brought by rivers. The soil formed due to weathering of limestone is regur soil and that of river is sticky black clayey soil.

Mineral potentiality based on geology, geophysics, ground geochemistry etc.

It was noted that the Milliolite limestone is the only rock formation which is exposed as outcrops in the entire sea-coast and especially in the area of investigation. It is exposed on the surface in forms of low mounds.

The typical characteristic of this limestone formation is its erosional feature. Due to chemical and differential weathering, solution holes, channels, cavities with alternate groove and ridge formation giving the area honeycomb appearance. This topography is known as Karst topography.

The limestone found is mostly light coloured, mostly white with shades of grey, buff, brown and yellow. The presence of impurities mostly iron is responsible for this colour variation. The rock is hard and compact. The texture of rock is mostly fine to medium grained at times ranging into coarse grade also. At places false and current beddings are also observed. In this limestone deposit the thickness is variable due to its topography.

Scope for proposed exploration

- 1. Location: Village: Lodhva Tehsil: Sutrapada, District: Gir Somnath, State: Gujarat.
- 2. Quantum of work: The approximate core drilling work is 360 meters.
- Rock formations to be drilled: Various types of soil, sub-soil, Limestone, Marl, clay, etc.
- The boreholes shall be in depth range of approximately 20 m. However, this is indicative only and may vary as per actual geological conditions.
- 5. Type of Drilling: Core drilling by Hydraulic Drilling Rigs.
- 6. Borehole size: The holes shall be derived in NQ sizes.
- The core recovery in all the formation should be at least 90% except in fault zone, weathered zone, soil, sand and structurally disturbed area
- CGM will provide proposed borehole location plan of all the areas to be covered under exploration.

9. Sampling: There will be 1 one mineral/rock sample for each 1-meter run of economically viable mineral. Each sample should be cut by core splitter. Each run shall be marked properly by plastic cards and the core boxes shall be numbered properly. Each sample shall be reduced to an approximate quantity by following the standard sampling procedures such as homogenizing, coning, quartering and pulverizing into 100/200 mesh and be prepared into two packets of 100-200 gm each. The final sample pockets shall be properly labelled with BH number, sample run.

Recommendations of G4 Stage Mineral Exploration Report.

A. K. Trivedi, 1979-81 concluded that the milliolite limestones of Saurashtra coasted area are most expensive deposits in Gujarat state. There deposits are the purest type of carbonate deposits. The area of they investigated has proved itself to be of very high potentiality with reserve of 138.24 million tons of limestone within an area of 4.8 Sq. Kms. The quality of limestone is very good. It is recommended that to continue the investigation to explore and study entire milliolite limestone of coastal area of Saurashtra. This study will establish its industrial utility for much more new industries in this part of the state.

U. D. G. Rao, 1965-66 had recommended that by virtue of Limestone deposit occurrence in the form of low ridges and mounds, these limestone deposits are eminently suited for economic exploration, by the simplest methods of quarrying. Other factors such as roads, transport facilities, availability of water and labour etc. present and can be arranged within reasonable costs, for the commercial exploration of any group of patches in the region examined.

S.D. Kapse, Y. C. Patel, 1997 also recommended that the study of the surveyed area revealed that limestone of Gaj and Dwarka formations are continuing to occur in adjacent sheets along-with laterite patches. So it is recommended to cover these areas under Pre-detailed survey.

Objectives of Exploration

- To know the continuity of the mineral body both along the strike and dip.
- · To map the extent of the ore body and lithology of the area.
- To ascertain the grade of Limestone deposit.
- Ore resource/reserve estimation in accordance with MEMC Rule-2015

2. Previous Work

Previous Exploration in proposed block area as well as adjoining area: All the sample (bed rock/trench/ groove/soil), borehole location should be plotted on the geological map and analytical data should be discussed briefly

S.D. Kapse, Y. C. Patel (Field Seson -1995-96-97) has done part of the Junagadh and Jamnagar districts of Saurashtra were taken up for geological mapping on 1:50,000 scales covering an area about 2580 sq. kms. by using Remote Sensing Techniques. The geological maps were prepared using False Colour Composite (F.C.C) OF "LANDSAT" and "SPOT" Satellite data. This area falls under the Survey of India topographical map 41 G/5, 41 G/10, 41 G/6, 41 K/4, 41G/15, 41 G/11, 41 L/5, 41 L/1, 41 L/9, 41 L/10.

U.D.G.Rao (Field Season-1965-66) has done the work eastern parts of Veraval and kodinar in the Gir Somnath District, Gujarat State. During the year 1965-66, was part of a project to assess the quality and quantity of Limestone for its economical utilization. During this field Season an area of about 200 sq.kms. between Kodinar and veraval was examined and all important limestone outcrops mapped and assessed And 228 chip sample was been collected systematically along and across the dip direction and sent for analysis. An area was found to be under limestone and the reserves have been estimated to be about 30 million tonnes of limestone and 26 million tonnes of calcareous sand.

A. K.Trivedi (Field Season 1979-80-81) has done the From Morasa to Padruka village of Veraval Taluka of Junagadh District, Gujarat state. During this Field Season an area Geological mapping on 1:2000 and contouring at 2mts interval were done covering an area of 4.8 SQ. Kms and total 36 boreholes with total meterage of 682.10 mts were drilled. The bore holes were located at 400 mts interval on a grid pattern of N-S, E-W. Except one or two most of the bore holes are drilled below 15 mts. with minimum depth 10.67 mts. and maximum depth of 30.50 mts. The deepest borehole of 30.50 mts was drilled to touch the trap rock. But inspite of this depth the trap could not be touched. The boreholes are mostly closed in gaj clays. The thickness of milliolite limestone strata varies from 6 mts to 21 mts. On an average the limestone thickness is 15 mts. The Total 654 sample was collected and 86 chip sample sent for analysis. An area was found to be under limestone and the reserves have been estimated to be about 138.24 million tonnes of cement grade limestone.

Commissioner of Geology and Mining (CGM), Gujarat had done on-ground G2 level exploration of limestone in the Barvela-Singhsar Block, Junagadh District. The exploration was carried out by drilling 53 boreholes at nearly 200-meter grid spacing. Drilling for Limestone in this region began in 2019 and was finished in 2021. A total of 51 boreholes tested positive for Limestone. All boreholes drilled in 50mts depth and still mineralized zone is continuing in the depth.

Ambuja Cement mining lease is 800 meters in west direction away from proposed block. Gujarat Sidhee Cement mining lease is 1200 meters in east direction away from proposed block. So, these lease areas are surrounding areas of the proposed block which testify the limestone prospectivity of this block.

Previously Geological survey of india conducted a geological mapping for Limestone, revealing the geological potential for Limestone in the area.

CGM/ GMRDS officials has also carried out field in current year survey of the proposed area and taken grab samples 08 Nos and done the chemical analysis of that samples. So chemical analysis also suggests that good quality limestone present in the study area, it was recommended to more sampling of the area. Chemical analysis data is given as per Annexure-1.

3. Block description

Block corner points	Latitude	Longitude		
1	20°51'13.08"N	70°35'6.65"E		
2	20°51'16.47"N	70°35'58.28"E		
3	20°50'33.01"N	70°36'40.89"E		
4	20°50'29.12"N	70°35'1.70"E		

4. Planned Methodology

- Preparation of Geological map. Field traverse and collection of grab/ channel samples from outcrop/ river cutting/ old dug wells.
- To carry out DGPS survey and identified location of Road, River, Settlement, Electric line, Telephone line or any other permanent structure.
- 3. Preparation of Landuse, Landpattern map along with ownership details.
- Preparation of proposed borehole location plan based on the previous work, grab sampling report and feasibility of drilling.
- Carry out systematic & scientific core drilling activity along with preparation of Lithounits, core drilling registers, sampling and preparation of samples for further physical, chemical and petrographic analysis.
- Carry out physical, chemical and petrographic analysis and interpretation of their result in borehole lithology.
- Preparation of draft Geological report as per specified format of NMET as per MEMC Rules, 2015 & its amendments.
- Submission of draft Geological report to designated Geological expert for peer review.
- Submission of final Geological Report incorporating comments of peer review to NMET for final approval.
- 10. Submission of Geological reports to state auction cell for further auction process.

5. Nature Quantum and Target

Components	G2				
Aerial reconnaissance	NA				
Geological Survey	NA NA				
Systematic drilling	Refer below (Plate-2)				
Petrographic and mineral graphic studies	Specific Gravity studies				
Synthesis of all available data	i) Integration of regional/ detailed geophysical, geological and geochemical data, if not done earlier.				
	ii) Synthesis of all available data and Report writing				

Borehole spacing (As per MEMC, 2015)

Type of deposit	Tabular deposit of regular	Bedded stratiform and tabular deposits of irregular habit (Minerals to be identified)	occurring en echelon				
G2 Stage	400 m	Not applicable	Not applicable				
	(Vertical depth of intersection of mineralised zone for different level boreholes should be specified, number of boreholes (first, second, third), borehole spacing, approximate length of different level of boreholes may also be specified)						

6. Exploratory Drilling

- The boreholes shall be in depth range of approximately 20 m. However, this is indicative only and may vary as per actual geological conditions.
- · Type of Drilling: Core drilling by Hydraulic Drilling Rigs.
- · Borehole size: The holes shall be derived in in NQ sizes.
- While drilling, wherever water table is encountered, depth of the water table should be recorded and to be mentioned in the driller logs.
- The core recovery in all the formation should be at least 90% except in fault zone, weathered zone, soil, sand and structurally disturbed area.

7. Manpower deployment

SI.	Activities	Unit			MO	NTHS		
No.	Activides	Unit	1	2	3	4	5	6
1	Camp Setting	Month						
2	Surface Drilling	m.						
3	Survey Party days (1 Party)	day						
4	Geologist Party days in field (1 Party)	day						
5	Sampling Party days, Core Sampling (1 party)	day						
6	Laboratory Studies	Nos.						
7	Camp Winding	Month						
8	Geologist Party days in HQ (1 Party)	day						
9	Geological Report Writing with Peer Review	Month						

Note: 1. Commencement of project may be reckoned from the day the exploration acreage is available along with all statutory clearances.

2. Time loss on account of monsoon/agricultural activity/forest clearance/local law & order problem may be additional to above time line.

8. Break-up of expenditure

The cost has been estimated based on actual schedule of rates mandated in the circularOM No. 61/1/2018/NMET dated 31st March 2020 for NMET funded projects which is **Rs. 88.93 Lakhs**. The detailed cost sheet for G-2 exploration for Limestone in proposed Lodhva Block is given below:

SL. NO.	Item	Estimated Cost (Rs.)		
1	Drilling	2439000		
2	Geology and Survey	1523520		
3	Laboratory	1794170		
	Sub Total (1 to 3)	5756690		
4	Miscellaneous	1779631		
	Total	7536321		
	GST 18%	1356538		
	Grand Total (including GST)	8892858		
	Say Rs. In Lakhs	88.93 Lakhs		

9. References

- "Geological report of southern part of Saurashtra covered under Toposheet Nos. 41 G/5, 41 G/10, 41 G/6, 41 K/4, 41G/15, 41 G/11, 41 L/5, 41 L/1, 41 L/9, 41 L/10 (Based on Interpretation of Satellite Imagery with Limited Field Checks)" (Field Season 1995-96 and 1996-97) by S.D. Kapse, Y. C. Patel, Commissioner of Geology and Mining, 1997.
- "Report of the Limestone Deposits in the Eastern parts of Veraval taluka" by U. D.
 G. Rao, Senior Geologist, Directorate of Geology and Mining, Ahmedabad, 1965-66.
- "A Report on the Limestone Drilling Scheme at Morasa Village of Veraval taluka of Junagadh District" (Field Season 1979-80-81) by A. K. Trivedi, Assistant Geologist, Directorate of Geology & Mining, Rajkot, 1981.
- "Preliminary Appraisal of Limestone in Coastal areas of Saurashtra, Gujarat for Selection of targets for Limestone suitable for S.M.S., B.F. and Chemical grade purposes" (Field Season 1997-98) by Dr. Manas Chowdhury, Jr. Geologist, Geological Survey of India.
- "Report on the Geological mapping in parts of toposheet nos. 41 L/9, L 13, K/12 and 41 P/5, Junagadh & Amreli District, Gujarat" (Field Season 1983-84) by A.C. Banerjee and A. Mukhopadhyay, Jr. Geologist, Geological Survey of India, 1985.

List of Annexures

Annexure-1: Chemical analysis data of collected samples.

Annexure-2: Detailed Cost Estimation Sheet

List of Plates

Plate 1: Proposed block boundary over existing Geological map.

Plate 2: Proposed Borehole Location Map.

Plate 3: Proposed block boundary over topographic map.

Plate 4: Satellite image of the proposed block showing surrounding existing Limestone leases


Block Id: CGM/NMET/Limestone/06/2024-25 Commissioner of Geology and Mining, Gujarat

Annexure-1


Cr No	Comple ID	Location		Loca	tion	SIO2	AL203	Fe2o3	TiO2	CaO	MgO	Na2O	K20	P2O5	503	LOI	TOTAL
Sr. No	Sample ID	Latitude	Longitude	%	96	%	- %	%	96	96	%	96	%	96	%		
1	LODHWA-01	20.847588*	70.58654°	7.67	4.21	3.28	0.45	47.87	0.85	0.00	0.29	0.10	0.05	34.88	99.65		
2	LODHWA-02	20°50'56.231"	70°35'18.590"	4.50	2.34	2.38	0.25	51.04	0.50	0.10	0.21	0.05	0.00	38.46	99.83		
3	LODHWA-03	20°50'50.69"	70°35'17.93"	3.59	1.79	2.69	0.26	53.11	0.67	0.14	0.18	0.13	0.00	37.24	99:80		
4	LODHWA-04	20°50'59.409"	70°35'55.367"	4.58	1.97	2.54	0.31	51.80	0.58	0.13	0.21	0.06	0.00	37.67	99.85		
5	LODHWA-05	20°51'02.364"	70°35'48.279"	3.50	1.55	2.45	0.22	52.83	0.51	0.14	0.15	0.50	0.00	38.00	99.85		
6	LODHWA-06	20*51'07.213"	70"35"53.926"	4.96	1.89	2.57	0.36	53.43	0.64	0.08	0.25	0.09	0.04	35.52	99.83		
7.	LODHWA-07	20*50'46.18"	70"36'14.71"	3.85	1.60	2.15	0.20	53.19	0.52	0.16	0.24	0.02	0.00	37.90	99.83		
8	LODHWA-08	20"50'40.13"	70"36'33.09"	5.62	1.53	2.25	0.28	52.16	0.35	0.08	0.08	0.14	0.03	37.29	99.81		

			Rates as per	NMET SoC	Estima	sted Cost of the
Sr. No.	Item of Work	Unit		2020-21		Proposa
21-110	TELLIO WORK	0.1116	SoC- Item	Rates as	000	Total Amoun
			S.No.	per SoC	Qty.	(Rs.
Α	DRILLING					,,,,,,
1	Surface Drilling	m.	2.2.1.1b	6775	360	243900
	Sub Total A					243900
В	Geological Work			1		
1	Survey Days	day	1.6.1a	8300	20	16600
	Labour (3 labour for survey)	day	5.7	504	60	3024
2	Geologist (Field)	day	1.3b	11000	30	33000
3	Geologist (HQ)	day		9000	40	36000
4	Core Sampling Days	day	1.5.2	5100	20	10200
	Labour (3 labour)	day	5.7	504	70	3528
5	Drilling camp setting/ winding cost	per drill	2.2.9a & 2.2.9b	250000	2	50000
	Sub-Total B					152352
C	LABORATORY STUDIES		1	T		
a	Chemical Analysis					
1	Primary Analysis	Nos.	4.1.15a	4200	360	151200
2	External Check Samples	Nos.	4.1.15a	4200	36	15120
В	Physical Analysis	10000	ANIMOTE ANI			250,000
1	Preparation of Thin Section	Nos.	4.3.1	2353	10	2353
2	Petrographic Studies	Nos.	4.3.4	4232	10	4232
3	Digital Micro Photographs	Nos.	4.3.7	280	5	140
4	Specific Gravity	Nos.	4.1	3540	18	6372
0.00	Sub-Total C	11027	200	22.0		179417
	Total (A+B+C)	Г Т	1	7		575669
D	Miscellaneous Charges	1	-	_	_	373003
1	Geological Report - 5	16	5.2			28783
2	Proposal Preparation- 2% of a project cost or 3.8 lakh (whice lower)	XD2549053755439	5.1			13293
3	Drill Core Preservation	m.	5.3	1590	360	57240
4	Peer Review Charges		1	30000		3000
5	Tender Process					13293
6	Operational Charges					62351
		Total		-		753632
	GST 18%					135653
	GRAND TOTAL					889285
	Rs. In Lakhs					88.93 lakh

Geological map of Lodhva Limestone Block, Gir-Somnath District, Gujarat



Block Id: CGM/NMET/Limestone/06/2024-25 Commissioner of Geology and Mining, Gujarat Plate - 1



Lodhva Limestone Block, Gir-Somnath District, Gujarat 70°34'30"E 70°36'0"E 70°37'30"E 20°51'0"N







Plate - 3

